

BEST AVAILABLE COPY

PCT/IB 04 / 01813

(06.07.04)

PA 1179641

06 JUL 2004

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

June 07, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 10/452,926

FILING DATE: June 03, 2003

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



T. LAWRENCE
Certifying Officer

Atty. Dkt. No. 023234-0103

15915 U.S. PTO
06/03/03**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Applicant: Marco FILICORI
Title: UNITARY COMBINATIONS OF FSH AND hCG
Appl. No.: Unassigned
Filing Date: 06/03/2003
Examiner: Unassigned
Art Unit: Unassigned

UTILITY PATENT APPLICATION
TRANSMITTAL

Mail Stop PATENT APPLICATION
Commissioner for Patents
PO Box 1450
Alexandria, Virginia 22313-1450

Sir:

Transmitted herewith for filing under 37 C.F.R. § 1.53(b) is the nonprovisional utility patent application of:

Marco Filicori

Enclosed are:

- [X] Application Data Sheet (37 CFR 1.76) (2 pages).
- [X] Unexecuted Declaration and Power of Attorney (3 pages).
- [X] Specification, Claims and Abstract (30 pages).
- [X] Formal drawings (2 sheets, Figures 1A, 1B and 2).

Atty. Dkt. No. 023234-0103

The filing fee is calculated below:

	Claims as Filed	Included in Basic Fee	Extra Claims	Rate	Fee Totals
Basic Fee				\$750.00	\$750.00
Total Claims:	31	- 20	= 11	x \$18.00	= \$198.00
Independents:	2	- 3	= 0	x \$84.00	= \$0.00
If any Multiple Dependent Claim(s) present:				+ \$280.00	= \$0.00
Surcharge under 37 CFR 1.16(e) for late filing of Executed Declaration and late payment of filing fee				+ \$130.00	= \$130.00
				SUBTOTAL:	= \$1078.00
[] Small Entity Fees Apply (subtract 1/2 of above):					= \$0.00
				TOTAL FILING FEE:	= \$1,078.00

- [] A check in the amount of \$0.00 to cover the filing fee is enclosed.
- [X] The required filing fees are not enclosed but will be submitted in response to the Notice to File Missing Parts of Application.
- [] The Commissioner is hereby authorized to charge any additional fees which may be required regarding this application under 37 C.F.R. §§ 1.16-1.17, or credit any overpayment, to Deposit Account No. 19-0741. Should no proper payment be enclosed herewith, as by a check being in the wrong amount, unsigned, post-dated, otherwise improper or informal or even entirely missing, the Commissioner is authorized to charge the unpaid amount to Deposit Account No. 19-0741.

Please direct all correspondence to the undersigned attorney or agent at the address indicated below.

Respectfully submitted,

Date 3 June 2003By S. A. Bent

FOLEY & LARDNER
Customer Number: 22428



22428

PATENT TRADEMARK OFFICE

Telephone: (202) 672-5404
Facsimile: (202) 672-5399

Stephen A. Bent
Attorney for Applicant
Registration No. 29,768

Application Data Sheet**Application Information**

Application Type:: Regular
Subject Matter:: Utility
Suggested classification::
Suggested Group Art Unit::
CD-ROM or CD-R?:: None
Computer Readable Form (CRF)?:: No
Title:: UNITARY COMBINATIONS OF FSH AND
hCG
Attorney Docket Number:: 023234-0103
Request for Early Publication?:: No
Request for Non-Publication?:: No
Suggested Drawing Figure:: 1
Total Drawing Sheets:: 2
Small Entity?:: No
Petition included?:: No
Secrecy Order in Parent Appl.?:: No

Applicant Information

Applicant Authority Type:: Inventor
Primary Citizenship Country:: Italy
Status:: Full Capacity
Given Name:: Marco
Family Name:: Filicori
City of Residence::
Country of Residence:: Bologna
Street of mailing address:: Vicolo Malgrado 2/8
40125 Bologna, Italy
Country of mailing address:: Italy

Correspondence Customer Number:: 22428

E-Mail address:: sbent@foleylaw.com

Representative Information

Representative Customer Number::	22428	
---	-------	--

Domestic Priority Information

Application::	Continuity Type::	Parent Application::	Parent Filing Date::

Foreign Priority Information

Country::	Application number::	Filing Date::	Priority Claimed::

Assignee Information

Assignee name::

10452926 .060303

Atty. Dkt. No.: 023234-0103

U.S. PATENT APPLICATION
for
UNITARY COMBINATIONS OF FSH AND hCG

Inventors: Marco Filicori

UNITARY COMBINATION OF FSH AND hCG

FIELD OF THE INVENTION

[0001] The present invention relates to novel gonadotropin formulations. More specifically, the invention relates to pharmaceutical formulations, useful for ovarian stimulation, in which both follicle stimulating hormone (FSH) and human chorionic gonadotropin (hCG) are present.

BACKGROUND OF THE INVENTION

[0002] Assisted reproduction technology (ART) procedures typically require treatment with exogenous gonadotropins to stimulate growth and maturation of the ovarian follicles. When gonadotropins are used to treat anovulatory females, the goal is to replicate the normal menstrual cycle, when a single, dominant follicle matures prior to induction of ovulation. In contrast, for women undergoing *in vitro* fertilization (IVF), controlled ovarian stimulation (COS) is employed to stimulate the growth and maturation of several ovarian follicles, yielding multiple oocytes, which then are retrieved for use in the IVF procedure.

[0003] Despite recent advances in ART, ovarian stimulation through exogenous gonadotropins is not uniformly successful due, in part, to varying individual responses to treatment with gonadotropins. This variability complicates patient management and can result in multiple births and potentially life-threatening complications.

[0004] Gonadotropins are secreted by the pituitary gland under the control of hypothalamic gonadotropin-releasing hormone (GnRH). Follicle stimulating hormone (FSH) and luteinizing hormone (LH) are the pituitary

hormones essential for follicular maturation (folliculogenesis) and luteinization. FSH is required for follicular recruitment (*i.e.*, the early growth of ovarian follicles) at the beginning of the spontaneous menstrual cycle, and it also supports mid- and late-stage folliculogenesis.

[0005] FSH is administered therapeutically to induce folliculogenesis in anovulatory women and women undergoing COS. In traditional ovulatory stimulation methods, FSH is administered throughout treatment until the time that oocytes are retrieved. This continued stimulation by FSH can cause multiple conceptions and the potentially fatal condition, ovarian hyperstimulation syndrome (OHSS). Decreasing the dosage of FSH can reduce the risk of OHSS, but low FSH dosages yield inadequate follicle quantities and thus lower the chances of success in assisted reproduction.

[0006] LH functions during all stages of a normal menstrual cycle. LH stimulates the theca cells of the follicle to produce the androgen substrate which is converted into estrogen by the aromatase system in the granulosa cells. During the late stages of follicle maturation, approximately 5 to 7 days before ovulation, large ovarian follicles begin to express LH receptors in granulosa cells, which render those follicles responsive to LH for continued maturation and development. Hillier *et al.*, *Mol. Cell Endocrinol.* 100:51 (1994), Campbell *et al. J. Reprod. Fertil.* 117:244 (1999). Next, a mid-cycle surge of LH triggers the final stage of follicular maturation and ovulation in a normal menstrual cycle. Ovulation follows the mid-cycle LH surge within 24 to 48 hours. Finally, in the second part of the menstrual cycle, the luteal phase, LH stimulated production of estrogen and progesterone in the corpus luteum of the ovary prepares the uterus for implantation and pregnancy.

[0007] In ovarian stimulation protocols, hCG can serve as a source of LH activity because hCG and LH act through the same receptor. Filicori

Atty. Dkt. No.: 023234-0103

et al. Human Reprod. 17:2009 (2002a); Martin *et al., Fertil. Steril.* 76: O-49 (2002). Relative to LH, hCG has a longer half-life and, hence, is more potent than LH, although the literature tends to treat hCG and LH as fungible. Indeed, the scientific literature generally does not mention determining the source of LH activity in naturally-derived gonadotropin preparations. *But see* Filicori *et al., Human Reprod. Update* 8: 543, 552 (2002b) ("likely hCG content of [a particular] hMG preparation" extrapolated to be "~ 5 IU per ampoule," such that, "of the 75 IU of LH potency contained in this hMG preparation, about 30 IU are provided by hCG").

[0008] The literature discloses using LH activity or low doses of hCG in combination with FSH throughout ovulatory stimulation, but guidance regarding effective amounts and timing of LH activity supplementation is lacking. For example, the abstract of Martin *et al., Fertil. Steril.* 76: O-49 (2002), discloses administering 2.5 µg recombinant hCG daily (maintaining serum hCG levels from 1-3 mIU/mL) during ovulatory stimulation. Gordon *et al.* disclose administering 75 IU FSH with 0, 1, 25, and 75 IU LH activity. *Human Reprod.* 12 (Suppl. 1): 52 (1997a); *ibid.*: 53 (1997b).

[0009] Published studies disclose administering LH activity, throughout stimulation, at FSH to LH ratios of 150:0, 150:37.5, 150:75, and 150:150. Filicori *et al.* (2002a). Further, the literature documents supplementing FSH stimulation with 50 IU hCG /day (Filicori *et al., J. Clin. Endocrinol. & Metabol.* 84: 2659 (1999)), and protocols in which 150 IU FSH is administered for 7 days, followed by treatment with FSH-to-hCG ratios of 150:0, 50:50, 25:100, and 0:200 (*ibid.* 87:1156 (2002c)).

[0010] The literature documents other compositions that contain both FSH and LH activity, as well as use of FSH in combination with LH activity. For example, PCT application WO 00/67778, published

Atty. Dkt. No.: 023234-0103

November 16, 2000, is directed to using LH or an equivalent amount of hCG in combination with FSH to induce folliculogenesis in anovulatory women. More particularly, the '778 application discloses administering LH or "a biologically-active analogue thereof" in doses of 100 to 1500 IU per day (page 4, lines 26-29) and in FSH:LH ratios that range from 1:1.5 to 1:20 (*id.*, lines 16-18).

[0011] United States patent No. 5,929,028 is directed to liquid formulations that contain one or more natural or recombinant gonadotropins, including FSH, LH, and hCG. The '028 patent discusses naturally derived compositions of human menopausal gonadotropin (hMG), which have FSH and LH activities in a ratio of approximately 1:1, but mentions no ratio of FSH to LH activity other than the 1:1 ratio of commercial hMG preparations.

[0012] Additionally, there are commercial formulations that contain both FSH and LH. Human-derived preparations are available containing 75 IU FSH with 75 IU LH activity (Pergonal, Humegon, Menogon, Repronex, and Menopur) and 75 IU FSH with 25 or 35 IU LH activity (Normegon and Pergogreen).

[0013] It is conventional wisdom, however, that "excessive" LH levels, albeit ill-defined, result in follicular atresia, suppression of granulosa cell proliferation, and premature luteinization. *See, generally, Filicori, Fertil. Steril.* 79: 253 (2003). Although recent work suggests otherwise, a notion persists in the field that LH activity levels must be within a certain range, and that levels below or above an "LH ceiling" impair normal follicle development. Shoham, *Fertil. Steril.* 70: 1170 (2002).

[0014] In summary, there is published evidence that supplementing FSH with LH activity during ovulation induction reduces the duration of treatment and the amount of gonadotropin used to achieve proper follicle development. Filicori *et al.* (1999), (2002b). On the other hand, the

belief persists that "high" LH activity levels negatively impacts follicle development.

[0015] That belief has guided the conventional ovarian-stimulation paradigm, which involves administration of FSH throughout controlled ovarian stimulation. Exogenous LH activity is deemed unnecessary and even detrimental during the early to middle stages of follicular development. Accordingly, the traditional means of ovarian stimulation entails treatment with FSH alone, typically at 75 IU/day. In this traditional protocol, LH activity is administered to induce ovulation only after the follicle reaches a certain stage of development. Only recently has LH activity been administered throughout treatment, and the optimal amount and timing of LH activity that is effective in this context remains controversial.

SUMMARY OF THE INVENTION

[0016] Conventional protocols of ovulation stimulation do not accommodate selective treatment regimens that optimize ovarian follicle development. Furthermore, available gonadotropic preparations are not easily adapted to different therapeutic goals. Thus, a need exists for compositions that combine FSH and hCG in varying ratios, thereby to enable the practitioner to tailor a gonadotropin treatment regimen to the needs of the individual patient. Methodology also is needed for using such compositions to stimulate folliculogenesis in anovulatory females and in the context of ART procedures.

[0017] To these ends, *inter alia*, the present invention provides pharmaceutical compositions that contain various amounts of FSH and hCG, as well as various FSH-to-hCG ratios, supplied in a single preparation. These compounds enable the practitioner to optimize

ovulatory stimulation in flexible manner not possible with available preparations.

[0018] The present invention also comprehends an approach to inducing ovulation, by using compositions with varying FSH:hCG ratios. The inventive methodology allows for incremental adjustments in the ratio of FSH to hCG, as a function of the stage of folliculogenesis or of variation in patient response, resulting in safer and more successful ovulatory stimulation.

[0019] One embodiment of the invention is a pharmaceutical composition that consists essentially of FSH and hCG in a pharmaceutically acceptable carrier. According to the invention, the ratio of FSH to hCG in such a composition is conducive, upon administration of the composition, to folliculogenesis and to follicular maturation without ovarian hyperstimulation. In this regard, use of the phrase "consists essentially of" means that the composition, while possibly having other constituents, does not include any component that inhibits or otherwise detracts from the beneficial properties of the composition with respect to folliculogenesis or follicular maturation.

[0020] Another embodiment of the invention is an assemblage comprising a first vial and a second vial, each of said vials containing a pharmaceutical composition consisting essentially of FSH and hCG in a pharmaceutically acceptable carrier, wherein the ratio of FSH to hCG is conducive, upon administration of said composition, to folliculogenesis and follicular maturation without ovarian hyperstimulation. The ratio of FSH to hCG differs between the first vial and the second vial.

[0021] A further embodiment of the invention is a method of inducing ovulation, comprising: administering at least one pharmaceutical composition characterized by a ratio of FSH to human hCG that is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5

Atty. Dkt. No.: 023234-0103

IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU
 hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU
 hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG,
 75 IU FSH:10 IU hCG, 75 IU FSH:25 IU hCG, 75 IU FSH:50 IU hCG, 75
 IU FSH:75 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU
 FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU
 FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU
 FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU
 FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150
 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU
 FSH:25 IU hCG, 150 IU FSH:75 IU hCG, 150 IU FSH:100 IU hCG, 150 IU
 FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200
 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU
 FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU
 FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and
 200 IU FSH:400 IU hCG, monitoring serum hormone levels, follicle size
 and follicle number; and then inducing ovulation by administration of an
 hCG bolus.

[0022] Other features, objects, and advantages of the present invention
 are apparent in the detailed description that follows. It should be
 understood, however, that the detailed description, while indicating
 preferred embodiments of the invention, are given by way of illustration
 only, not limitation. Various changes and modifications within the spirit
 and scope of the invention will become apparent to those skilled in the art
 from the detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] FIGURE 1 shows useful FSH-to-hCG ratio values, where FSH amount is expressed in international units (IU) and hCG amount is expressed in IU (Fig. 1A) and micrograms (Fig. 1B).

[0024] FIGURE 2 is a bar graph that depicts a proposed treatment protocol according to the invention, using FSH:hCG compositions as described above.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0025] The invention provides compositions and methods for a novel therapeutic paradigm, characterized by administration of hCG in combination with FSH during all stages of treatment and by an adjustment of the FSH-to-hCG ratio, to optimize ovulatory stimulation. This paradigm departs substantially from the conventional approach, where FSH is administered alone, typically at 75-300 IU/day, and a bolus of LH activity is administered mid-cycle to induce ovulation.

[0026] The invention provides pharmaceutical compositions with a wide range of FSH to hCG ratios. The compositions and methods of the invention enable the physician to easily tailor treatment to a given patient's situation, allowing the ratio of FSH to hCG to be fine-tuned based on different stages of folliculogenesis and varying patient response to gonadotropins. Further, having a range of therapeutic compositions available, including those with high levels of hCG, allows the physician to easily accommodate the therapeutic requirements of various ART procedures. This flexibility is not possible with current gonadotropic preparations.

[0027] Because the compositions of the invention have a broad range of FSH to hCG ratios, the invention can easily be used to stimulate folliculogenesis and ovulation in any ovulation induction and ART

Atty. Dkt. No.: 023234-0103

procedure including among others, treatment of anovulatory infertility, in-vitro fertilization (IVF), intracytoplasmic sperm injection (ICSI), gamete intrafallopian transfer (GIFT), zygote intrafallopian transfer (ZIFT), cryopreserved embryo transfer, intrauterine insemination (IUI), donor oocyte transfer, cryopreserved embryo transfer from donor oocytes, and ART cycles for host uterus transfer. The compositions and methods of the invention can also be used to treat anovulatory patients and patients with hypogonadotropic hypogonadism and the polycystic ovary syndrome.

[0028] The compositions of the invention employ hCG as a source of LH activity. hCG binds to LH receptors and exerts through them its biological actions; separate hCG receptors do not exist. Using hCG in accordance with the invention has significant advantages over the use of LH. For instance, human-derived hCG is less expensive than either human-derived or recombinant FSH. From a cost perspective, therefore, it is favorable to minimize the amount of FSH used during the stimulation protocol. Furthermore, the use of hCG in a pharmaceutical composition of the invention more physiologically and effectively supports the final stages of maturation of large and mature ovarian follicles during COS, increases ovarian and blood estrogen levels, improves oocyte and embryo quality, lowers the amount and duration of FSH administration, and reduces the risk of FSH-associated complications such as multiple conceptions and OHSS.

[0029] In summary, the compositions and methods of the invention empower treatment regimens that achieve higher rates of success, that decrease the risk of complications, that are easy to implement, and that are less costly than those of conventional practice.

[0030] Unless indicated otherwise, all technical and scientific terms are used in a manner that conforms to common technical usage. Generally, the nomenclature of this description and the described laboratory

Atty. Dkt. No.: 023234-0103

procedures are well known and commonly employed in the art. Standard techniques are used for recombinant nucleic acid methods, microbial culture, cell culture, tissue culture, transformation, analytical chemistry, organic synthetic chemistry, chemical syntheses, chemical analysis, and pharmaceutical formulation and delivery. Generally, enzymatic reactions and purification and/or isolation steps are performed according to the manufacturers' specifications. Absent an indication to the contrary, the techniques and procedures in question are performed according to conventional methodology disclosed, for example, in Sambrook *et al.*, MOLECULAR CLONING A LABORATORY MANUAL, 2d ed. (Cold Spring Harbor Laboratory Press, 1989), and CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1989).

Unitary Combinations of FSH and hCG

[0031] The invention provides novel pharmaceutical compositions useful for inducing ovarian stimulation in a mammal. "Mammal" refers to a human, non-human primate, sheep, pig, cow, horse, donkey, mouse, rat, rabbit, guinea pig, dog, cat, or captive wild animal. Preferably, the mammal is a human.

[0032] The pharmaceutical compositions contain 50 to 200 IU FSH together with 1 to 400 IU human-derived hCG (see FIGURE 1A).

Preferably, the compositions contain FSH and hCG in the following ratios:

50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:25 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:75 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU

Atty. Dkt. No.: 023234-0103

hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:75 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

[0033] More preferably, the compositions contain FSH and hCG in the following ratios: 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:75 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

[0034] In a more preferred embodiment, the composition contains 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, or 50 IU FSH:400 IU hCG.

These ratios are useful in treatment protocols requiring multiple

administrations because more than one ampoule can be easily administered.

[0035] In another, more preferred embodiment, the composition contains 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, or 100 IU FSH:400 IU hCG. These ratios allow a physician to select compositions with a higher dose of FSH. These compositions can be particularly useful at the outset of treatment where higher doses of FSH may be desired.

[0036] In a further more preferred embodiment, the composition contains 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, or 100 IU FSH:25 IU hCG. Such compositions provide lower doses of hCG which can be advantageous at the onset of treatment.

[0037] Purified FSH can be obtained by any methods known in the art. FSH, as used herein, includes human-derived and recombinant FSH, FSH analogs, as well as deglycosylated, unglycosylated, and modified glycosylated forms.

[0038] Human-derived FSH can be purified by any means known in the art from natural sources such as urine, pituitary, and placenta. Procedures for isolating human-derived FSH are described in, *e.g.*, Fevold *et al. Endocrinology* 26:999 (1940), Fraenkel-Conrat *et al., Proc. Soc. Exp. Biol. Med.* 45:627 (1940), McShan and Meyer, *J. Biol. Chem.* 135:473 (1940), Greep *et al., ibid.* 133:289 (1940), Li *et al., Science* 109:445 (1949), and Roos and Gemzell, CIBRA FOUNDATION STUDY GROUP, No. 22 (Little, Brown and Co., Boston, MA, 1965).

[0039] Purified human-derived FSH is commercially available and is sold under the names Fostimon® (AMSA/IBSA), Metrodin HP® (Serono), and Bravelle (Ferring). Recombinant FSH can be obtained by any of several known means. For example, Keene *et al., J. Biol. Chem.* 26:4769 (1989), and WO 86/04589 describe expression and purification of biologically active human FSH in rodent cells.

Atty. Dkt. No.: 023234-0103

[0040] Recombinant FSH is also commercially available under the names Follistim® (Organon), Puregon® (Organon), and Gonal-F® (Serono).

[0041] It can be advantageous to employ FSH isoforms that differ in the extent to which they are post-translationally modified. Due to different modifications, the isoforms exhibit differences in overall charge, degree of sialic acid (a terminal sugar) or sulfate incorporation, receptor binding affinity and plasma half-life. Chappel *et al.*, *Endocrine Reviews* 4:179 (1983); Snyder *et al.* *Mol. Cell. Endocrin.* 54:115 (1987). These forms are separable from each other on the basis of their overall charge and all isoforms exhibit biological activity. Isoforms that exhibit a greater net negative charge are more heavily sialylated, exhibit a longer metabolic clearance rate and a greater biologic activity due to their extended plasma survival time.

[0042] hCG can be obtained by any means known in the art. hCG, as used herein, includes human-derived and recombinant hCG. Human-derived hCG can be purified from any appropriate source (*e.g.* urine, and placenta) by any method known in the art as disclosed in Zondek and Aschheim, *Klin. Wochenschr.* 7: 931 (1928), Katzman *et al.*, *ibid.* 148:501 (1943), and Claesson *et al.*, *Acta Endocrinol.* 1: 1 (1948), among others. Purified human hCG is commercially available and is sold under the names Profasi HP® (Serono), Gonasi® (AMSA/IBSA), and Choragon, Novarelin (Ferring). Methods of expressing and purifying recombinant hCG are known in the art and are disclosed, for example, in Gupta and Dighe, *J. Mol. Endocrinology* 22: 273 (1999).

[0043] Recombinant hCG possesses approximately 25 times greater activity by weight as compared human-derived hCG. For example, see Chang *et al.*, *Fertil. Steril.* 76: 67 (2001); The European Recombinant Human Chorionic Gonadotrophin Study Group, *Fertil. Steril.* 75: 1111 (2001). Thus, the amount of hCG can be adjusted accordingly to provide

Atty. Dkt. No.: 023234-0103

the desired international units of hCG activity when recombinant hCG is used in the compositions. An adjustment along this line, based on the values of FIGURE 1A, is represented in FIGURE 1B.

[0044] The amount of FSH and hCG activity can be determined using assay methods known in the art. 1 IU of hCG is equivalent to 5-7 IU LH in the pharmacopaeia Van Hell bioassay. Van Hell *et al.*, *Acta Endocrin.* 47: 409 (1964). hCG activity in a composition can be determined using any assay methods known, including the Van Hell bioassay, radioreceptor assays as described, for instance, in Dighe & Moudgal. *Arch. Biochem. Biophys.* 225: 490 (1983), whole animal bioassays, ovarian ascorbic acid depletion assay, and the MA10 Leydig cell bioassay disclosed Ascoli, *Endocrinology* 108: 88 (1981). Similarly, FSH activity can be determined using receptor binding assays and whole animal bioassays.

[0045] The amount of FSH and hCG protein in a given composition can be determined by the weight of the solid compound, protein assays, such as Bradford and Lowry assays, and immunoassay techniques such as ELISA and Western blotting.

[0046] FSH:hCG compositions can be formulated by admixing, in an aqueous solution, purified FSH and hCG products, with batch-wise adjustments to achieve the desired FSH:hCG ratio, followed by sterile filtration, sterile filling, and, if desired, lyophilization. The FSH:hCG composition also can be formulated using a method, wherein purification conditions are established that yield the desired FSH:hCG ratio in the first instance (*i.e.*, without compounding as such).

[0047] The pharmaceutical compositions of the present invention can be formulated into well known compositions for any route of drug administration, *e.g.*, oral, rectal, parenteral (intravenous, intramuscular, or subcutaneous), intracisternal, intravaginal, intraperitoneal, local (powders, ointments or drops), or as a buccal or nasal spray. A typical composition

Atty. Dkt. No.: 023234-0103

for such purpose comprises a pharmaceutically acceptable carrier, such as aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described in REMINGTON'S PHARMACEUTICAL SCIENCES, 15th Ed. (Mack Publishing Co., 1975), at pages 1405-12 and 1461-87, and THE NATIONAL FORMULARY XIV, 14th Ed. (American Pharmaceutical Association, 1975), among others.

[0048] Examples of suitable aqueous and non-aqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate.

[0049] The compositions of the present invention also can contain additives such as but not limited to preservatives, wetting agents, emulsifying agents, and dispersing agents. Antibacterial and antifungal agents can be included to prevent growth of microbes and includes, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. Furthermore, it may be desirable to include isotonic agents such as sugars, sodium chloride, and the like.

[0050] In some cases, to effect prolonged action it is desirable to slow the absorption of FSH and hCG from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of FSH and hCG then depends upon its rate of dissolution which, in turn, can depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered FSH and hCG combination form is accomplished by dissolving or suspending the FSH and hCG combination in an oil vehicle.

[0051] Injectable depot forms can be made by forming microencapsule matrices of the FSH and hCG combination in biodegradable polymers such

as polylactide-polyglycolide. Depending upon the ratio of FSH and hCG combination to polymer and the nature of the particular polymer employed, the rate of FSH and hCG combination release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the FSH and hCG combination in liposomes or microemulsions which are compatible with body tissues.

[0052] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use. Injectable formulation can be supplied in any suitable container, *e.g.* vial, pre-filled syringe, injection cartridges, and the like.

[0053] The pH and exact concentration of the various components of the pharmaceutical composition are adjusted in accordance with routine practice in this field. See GOODMAN AND GILMAN'S THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS, 7th ed.

[0054] In a preferred embodiment, the compositions of the invention are supplied as compositions for parenteral administration. General methods for the preparation of the parenteral formulations are known in the art and are described in REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY, *supra*, at pages 780-820. The parenteral compositions can be supplied in liquid formulation or as a solid which will be mixed with a sterile injectible medium just prior to administration. In an especially preferred embodiment, the parenteral compositions are supplied in dosage unit form for ease of administration and uniformity of dosage.

[0055] "Dosage unit form" in this description refers to physically discrete units that are suited as unitary dosages for a mammalian subject to be treated, where each unit contains a predetermined quantity of active

Atty. Dkt. No.: 023234-0103

material calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active material and the particular therapeutic effect to be achieved and the limitations inherent in the art of compounding active materials for use in mammalian subjects.

[0056] A unit dosage form can contain, for example, 50, 75, 100, 150, and 200 IU FSH in combination with 1, 5, 10, 25, 50, 75, 100, 200, 300, and 400 IU hCG. Expressed in proportions, FSH and hCG each are generally present in from about 0.1 μ g to about 2000 mg/ml.

[0057] Suspensions can contain rheology modifying agents such as, *e.g.*, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

[0058] One of ordinary skill will appreciate that effective amounts of the FSH and hCG and the proper ratios of FSH to hCG can be determined empirically. The compositions can be administered to a subject, in need of ovarian stimulation, as pharmaceutical compositions in combination with one or more pharmaceutically acceptable excipients. It will be understood that, when administered to a human patient, the total daily usage of the agents or composition of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors: the type and degree response to be achieved; activity of the specific agent or composition employed; the specific agents or composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the agent;

Atty. Dkt. No.: 023234-0103

the duration of the treatment, and like factors well known in the medical arts.

**Use of "Single-Vial" Combinations of FSH and hCG
in Ovulatory Stimulation**

[0059] Based on the treatment goals, a composition with the proper FSH to hCG ratio can be selected and the composition or a series of compositions can be administered throughout the period of ovarian stimulation. The physician can initiate stimulation with one ratio and then adjust the ratios of FSH to hCG throughout the cycle. Adjustments made during treatment can be based on the timing of administration during the cycle or in response to folliculogenetic or steroidogenetic indicators, such as the number and size of follicles and patient hormone levels.

[0060] For example, in treating anovulatory females characterized by hypogonadotropic amenorrhea, compositions of the invention can be selected and administered to mimic the levels of FSH and LH activity that occur during a normal menstrual cycle. A composition with an FSH to hCG ratio of 100:5 initially would be administered daily. Starting on day six, for example, the levels of serum estradiol and follicle number and size would be monitored. Once at least two follicles greater than 12 mm and estradiol levels higher than 200 pg/mL are detected treatment will continue using a composition with an FSH to hCG ratio of 50:200 which will be administered until the final maturation parameters of at least one follicle greater than 17 mm and estradiol levels higher than 400 pg/mL are achieved. Upon reaching the final maturation parameters, ovulation will be triggered with 10,000 IU of hCG.

[0061] During COS, compositions are selected to maximize the number of large, mature follicles while minimizing the number of small, potentially dangerous follicles. A composition with an FSH to hCG ratio of 250:10 initially would be administered daily. Beginning on day six, the levels of

Atty. Dkt. No.: 023234-0103

serum estradiol and follicle number and size would be monitored. Once at least four follicles greater than 12 mm and estradiol levels higher than 600 pg/mL are detected a composition with an FSH to hCG ratio of 50:200 will be administered until the final maturation parameters of at least five follicles greater than 17 mm and estradiol levels higher than 1,500 pg/mL are achieved. Then, ovulation would be triggered with 10,000 IU of hCG.

[0062] The pharmaceutical compositions can be coadministered with one or more other compounds or molecules. "Coadministered" refers to simultaneous administration in the same formulation or in two different formulations.

[0063] The compositions of the invention can be administered with or following GnRH agonists and antagonists. GnRH agonists and antagonists are used for ovulation induction procedures to prevent spontaneous ovulation during gonadotropin administration. GnRH antagonist abolishes endogenous LH activity, resulting in fewer viable oocytes. Accordingly, the compositions of the invention can be administered at the commencement of treatment with GnRH antagonists to provide LH activity levels which are sufficient to stimulate follicle development and oocyte maturation.

[0064] To provide additional flexibility, the compositions of the invention can be coadministered with the currently available gonadotropin preparations, such as pure FSH, pure hCG, pure LH, and hMG preparations. For example, 1 ampoule of the FSH:hCG combination composition having FSH 75 IU and hCG 200 IU can be coadministered with 1 ampoule of highly purified FSH containing FSH 75 IU resulting administration of 150 IU of FSH and 200 IU of hCG.

[0065] FSH:hCG compositions are administered sequentially throughout ovarian stimulation. "Sequential" administration refers to a time

Atty. Dkt. No.: 023234-0103

difference of from seconds, minutes, hours or days between the administration of the compositions. Each subsequent sequential administration can be comprised of a composition with the same ratio of FSH to hCG as the previous administration or a different ratio of FSH to hCG. In a preferred embodiment, sequential administration is performed with FSH:hCG compositions that differ in the ratio of FSH to hCG. In a more preferred embodiment each succeeding composition contains a greater amount of hCG over the preceding composition in the series. In an especially preferred embodiment, the ratio of FSH to hCG is changed incrementally as shown in Figure 2.

[0066] In the most preferred embodiment, the ratio of FSH to hCG is adjusted throughout the series in response to patient reaction to the FSH:hCG composition as indicated by folliculogenetic and steroidogenetic markers. When 2-4 intermediate follicles (12-14 mm) are formed and estradiol levels of 200-600 pg/mL are achieved, this is an indication that adequate folliculogenesis has been achieved by an FSH-rich preparation and it is appropriate to switch to an hCG-rich preparation.

[0067] Folliculogenesis and steroidogenesis should be monitored throughout ovarian stimulation by any means known in the art. Monitoring hormone levels and follicle size provides information regarding ovarian response during the treatment regimen and allows the physician to adjust the ratio of FSH to hCG during treatment. Follicle size can be determined, for example, using transvaginal pelvic ultrasonography. Follicles are categorized as large (> 14 mm), intermediate (10-14 mm) or small (< 10 mm). Transvaginal pelvic ultrasound can be performed frequently, for example on treatment days 0, 6, 8, 10, 12, 14, 16, 18, and 20, until preovulatory hCG administration.

Atty. Dkt. No.: 023234-0103

[0068] The levels of estradiol can be measured using any means known in the art from any appropriate body fluid (*e.g.*, blood, urine, and saliva), using an immunoassay or a chemiluminescence assay.

[0069] The following example illustrates the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples.

Controlled Ovulatory Stimulation for In Vitro Fertilization

[0070] Treatment is initiated in the mid-luteal phase of a spontaneous menstrual cycle when a GnRH agonist is administered. Ovulation induction is started 1-2 weeks later, after spontaneous menses. Alternatively, a GnRH antagonist is administered starting on the 6th day of ovarian stimulation. Patients receive daily injections of 200 IU FSH:10 IU hCG from day 1 and continuing until the appearance of at least 4 follicles > 11 mm and serum estradiol levels of > 600 pg/mL. Starting on this day and until the end of treatment patients receive daily administration of 50 IU FSH:200 IU hCG.

[0071] Treatment monitoring is conducted throughout FSH:hCG administration. Each day one blood sample is drawn between 0800-0900 and serum samples are prepared in the standard fashion. Estradiol levels are determined. Transvaginal ultrasound is performed frequently during FSH:hCG treatment and until just prior to preovulatory hCG administration. Estradiol levels are monitored using chemiluminescence assays (Chiron Corp. Diagnostics ACS 180, Milan, Italy).

[0072] Upon achievement of final maturation parameters – four follicles of > 14 mm and 17 β -estradiol levels of 800-1500 pg/mL - a bolus of hCG, *e.g.* 10,000 IU hCG is administered to trigger ovulation and oocytes are retrieved.

Atty. Dkt. No.: 023234-0103

WHAT IS CLAIMED IS:

1. A pharmaceutical composition consisting essentially of FSH and hCG in a pharmaceutically acceptable carrier, wherein the ratio of FSH to hCG is conducive, upon administration of said composition, to folliculogenesis and follicular maturation without ovarian hyperstimulation.

2. The composition of claim 1 free from any other proteins of mammal origin.

3. The composition of claim 1 wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:25 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:75 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

4. The composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU

Atty. Dkt. No.: 023234-0103

hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:50 IU hCG, 150 IU FSH:75 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

5. The pharmaceutical composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:100 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, and 200 IU FSH:200 IU hCG.

6. The pharmaceutical composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:75 IU

Atty. Dkt. No.: 023234-0103

hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

7. The pharmaceutical composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, and 50 IU FSH:400 IU hCG.

8. The pharmaceutical composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, and 100 IU FSH:400 IU hCG.

9. The pharmaceutical composition of claim 3 wherein the ratio of FSH to hCG is selected from the group consisting of 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, and 100 IU FSH:25 IU hCG.

10. The pharmaceutical composition according to claim 1, wherein said FSH is human-derived FSH.

11. The pharmaceutical composition according to claim 1, in lyophilized form.

12. The pharmaceutical composition according to claim 1, in unit dosage form.

13. The pharmaceutical composition according to claim 12, in solid dosage form.

14. The pharmaceutical composition according to claim 13, wherein the solid dosage form is selected from the group consisting of capsules, tablets, suppositories, pills, powders, and granules.

Atty. Dkt. No.: 023234-0103

15. The pharmaceutical composition according to claim 1 in liquid form.

16. The pharmaceutical composition according to claim 15 wherein the liquid form is supplied in a vial.

17. The pharmaceutical composition according to claim 15 wherein the liquid form is supplied in a pre-filled syringe or cartridge.

18. An assemblage comprising a first vial and a second vial, each of said vials containing a pharmaceutical composition according to claim 1, wherein the ratio of FSH to hCG differs between the first vial and the second vial.

19. The assemblage according to claim 18 further comprising written instructions on the timing for administering the compositions contained in the first and second vials.

20. A method of inducing ovulation, comprising:

(A) administering at least one pharmaceutical composition characterized by a ratio of FSH to hCG that is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:25 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:75 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU

Atty. Dkt. No.: 023234-0103

FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

(B) monitoring serum hormone levels, follicle size and follicle number; and then

(C) inducing ovulation by administration of an hCG bolus.

21. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:1 IU hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:25 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:100 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, 200 IU FSH:200 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

22. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:1 IU hCG, 50 IU FSH:5 IU hCG, 50 IU FSH:10 IU hCG, 50 IU FSH:25 IU hCG, 75 IU FSH:1 IU

Atty. Dkt. No.: 023234-0103

hCG, 75 IU FSH:5 IU hCG, 75 IU FSH:10 IU hCG, 75 IU FSH:50 IU hCG, 75 IU FSH:100 IU hCG, 100 IU FSH:1 IU hCG, 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, 100 IU FSH:25 IU hCG, 100 IU FSH:50 IU hCG, 100 IU FSH:75 IU hCG, 100 IU FSH:100 IU hCG, 150 IU FSH:1 IU hCG, 150 IU FSH:5 IU hCG, 150 IU FSH:10 IU hCG, 150 IU FSH:25 IU hCG, 150 IU FSH:100 IU hCG, 150 IU FSH:200 IU hCG, 200 IU FSH:1 IU hCG, 200 IU FSH:5 IU hCG, 200 IU FSH:10 IU hCG, 200 IU FSH:25 IU hCG, 200 IU FSH:50 IU hCG, 200 IU FSH:75 IU hCG, 200 IU FSH:100 IU hCG, and 200 IU FSH:200 IU hCG.

23. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:75 IU hCG, 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, 50 IU FSH:300 IU hCG, 50 IU FSH:400 IU hCG, 75 IU FSH:200 IU hCG, 75 IU FSH:300 IU hCG, 75 IU FSH:400 IU hCG, 100 IU FSH:200 IU hCG, 100 IU FSH:300 IU hCG, 100 IU FSH:400 IU hCG, 150 IU FSH:300 IU hCG, 150 IU FSH:400 IU hCG, 200 IU FSH:300 IU hCG, and 200 IU FSH:400 IU hCG.

24. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 50 IU FSH:100 IU hCG, 50 IU FSH:200 IU hCG, and 50 IU FSH:400 IU hCG.

25. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 100 IU FSH:100 IU hCG, 100 IU FSH:200 IU hCG, and 100 IU FSH:400 IU hCG..

26. The method of claim 20, wherein the ratio of FSH to hCG is selected from the group consisting of 100 IU FSH:5 IU hCG, 100 IU FSH:10 IU hCG, and 100 IU FSH:25 IU hCG.

27. The method of claim 20, wherein step (A) comprises administering in series at least two pharmaceutical compositions,

characterized by a ratio of FSH to hCG selected from said group, that is either the same or differs with respect to said ratio.

28. The method of claim 20, wherein each succeeding composition in said series contains hCG that is increased over the preceding composition in said series.

29. The method of claim 27, wherein the period of time between compositions of the series is selected from the group consisting of from 1 hour, 5 hours, 10 hours, 12, hours, 24 hours, 1 day 2, days . 3, days 4, days, 5 days, 6, days, 7, days, 8 days, 9 days, 10 days, 11, days, 11 days, 12, days, 13, days, 14 days, and 15 days.

30. The method of claim 20, wherein the composition further comprises pure FSH.

31. The method of claim 20, wherein the composition further comprises pure hCG.

ABSTRACT OF THE DISCLOSURE

A novel ovulatory induction paradigm entails administration of hCG in combination with FSH during all stages of treatment, where the ratio of FSH to hCG is adjusted to optimize ovulatory stimulation and minimize complications. The use of compositions characterized by various FSH:hCG ratios enables the practitioner readily to tailor the treatment regimen and accommodate different therapeutic goals as well as individual patient responses to gonadotropin administration.

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I HEREBY DECLARE:

THAT my residence, post office address, and citizenship are as stated below next to my name;

THAT I believe I am the original, first, and sole inventor (if only one inventor is named below) or an original, first, and joint inventor (if plural inventors are named below or in an attached Declaration) of the subject matter which is claimed and for which a patent is sought on the invention entitled

UNITARY COMBINATIONS OF FSH AND hCG

(Attorney Docket No. 023234-0103)

the specification of which (check one)

 X is attached hereto.

 was filed on as United States Application Number or PCT International Application Number and was amended on (if applicable).

THAT I do not know and do not believe that the same invention was ever known or used by others in the United States of America, or was patented or described in any printed publication in any country, before I (we) invented it;

THAT I do not know and do not believe that the same invention was patented or described in any printed publication in any country, or in public use or on sale in the United States of America, for more than one year prior to the filing date of this United States application;

THAT I do not know and do not believe that the same invention was first patented or made the subject of an inventor's certificate that issued in any country foreign to the United States of America before the filing date of this United States application if the foreign application was filed by me (us), or by my (our) legal representatives or assigns, more than twelve months (six months for design patents) prior to the filing date of this United States application;

THAT I have reviewed and understand the contents of the above-identified specification, including the claim(s), as amended by any amendment specifically referred to above;

THAT I believe that the above-identified specification contains a written description of the invention, and of the manner and process of making and using it, in such full, clear, concise, and exact terms as to enable any person skilled in the art to which it pertains, or with which it is most nearly connected, to make and use the invention, and sets forth the best mode contemplated by me of carrying out the invention; and

THAT I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

Atty. Dkt. No. 023234-0103

I HEREBY CLAIM foreign priority benefits under Title 35, United States Code § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign application for patent or inventor's certificate or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number	Country	Foreign Filing Date	Priority Claimed?	Certified Copy Attached?

I HEREBY CLAIM the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below.

U.S. Provisional Application Number	Filing Date

I HEREBY CLAIM the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Application Number	Parent Filing Date	Parent Patent Number

I HEREBY APPOINT the registered attorneys and agents at Customer Number 22428.



22428

PATENT TRADEMARK OFFICE

to have full power to prosecute this application and any continuations, divisions, reissues, and reexaminations thereof, to receive the patent, and to transact all business in the United States Patent and Trademark Office connected therewith.

Atty. Dkt. No. 023234-0103

I request that all correspondence be directed to:

Stephen A. Bent
FOLEY & LARDNER
Customer Number: 22428



22428

PATENT TRADEMARK OFFICE

Telephone: (202) 672-5404
Facsimile: (202) 672-5399

I UNDERSTAND AND AGREE THAT the foregoing attorneys and agents appointed by me to prosecute this application do not personally represent me or my legal interests, but instead represent the interests of the legal owner(s) of the invention described in this application.

I FURTHER DECLARE THAT all statements made herein of my own knowledge are true, and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Name of first inventor	Marco Filicori
Residence	Bologna,
Citizenship	Italy
Post Office Address	Vicolo Malgrado 2/8 40125 Bologna, Italy
Inventor's signature	
Date	

Title: UNITARY COMBINATIONS

OF FSH AND hCG

Inventor(s): Marco FILICORI

DOCKET NO.: 023234-0103

10452926.060303

FIGURE 1A

	FSH (IU)				
	50	75	100	150	200
hCG (IU)	1	1	1	1	1
	5	5	5	5	5
	10	10	10	10	10
	25	25	25	25	25
	50	50	50	50	50
	75	75	75	75	75
	100	100	100	100	100
	200	200	200	200	200
	300	300	300	300	300
	400	400	400	400	400

FIGURE 1B


	FSH (IU)				
	50	75	100	150	200
Recombinant hCG (μ g)	0.04	0.04	0.04	0.04	0.04
	0.2	0.2	0.2	0.2	0.2
	0.4	0.4	0.4	0.4	0.4
	1	1	1	1	1
	2	2	2	2	2
	3	3	3	3	3
	4	4	4	4	4
	8	8	8	8	8
	12	12	12	12	12
	16	16	16	16	16

Title: UNITARY COMBINATIONS
OF FSH AND hCG

Inventor(s): Marco FILICORI
DOCKET NO.: 023234-0103

1452926 .060305

FIGURE 2



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.